



# LetCo

LIQUID ENERGY TRADING COMPANY SWISS S.A.

## Soybean Molasses

(Product code LQS-SBZ4S)

SOYBEAN MOLASSES is an important commercial and biological product, and a source of phytochemicals and soy sugars. The product is non- GMO.



### Description:

Color: dark brown  
Flavor: typical  
Odor: typical  
Bulk density (g/ml): 1,2 – 1,3  
Energy value: 730kJ(172kca)/100g  
Shelf life: 12 months  
Packing & Transport: in bulk

**Quality management:** Identity Preserved Program / Traceability

### Relevant identified uses:

Industrial source of soy phytochemicals (e.g. isoflavones) and soy sugars, feed ingredient, as a fermentation aid/substrate, as a prebiotic, as an ingredient in specialized breads, stabilizer, possible use as plywood adhesive

### Chemical Parameters

Parameters	Unit	Typical	Guarantee
Protein (N x 6,25), dry base (%)	%	11,0	max 12,0
Dry matter (%)	%	55,0	60,0 +/- 10
Fat, dry base (%)	%	1,0	max. 1,0
Carbohydrates (%)	%	45,0 by difference	

Nitrogen content is based on dry matter from 0,64 % to 1,3 % on the basis of statistical data.

### Microbiological Parameters

Total plate count	max. 12000000 / g
Salmonella	Negative in 25 g
Yeast and moulds	Max. 200000 / g

### Heavy Metals

Parameters	Unit	Typical	Guarantee
Lead, Pb	mg/kg	2,0	max.10
Cadmium, Cd	mg/kg	0,01	max 1,0
Mercury, Hg	mg/kg	0,001	max. 0,1
Arsenic, As	mg/kg	0,015	max. 2

### Periodical Analysis

Parameter	Unit	Guarantee
Fiber, dry base (%)	%	max. 1,0
Ash, dry base (%)	%	max. 12,0

### General information :

Soybean molasses is produced by aqueous ethanol extraction of white flakes and with addition of propionic acid and ammonium propionate.

Soya Molasses is a co-product of the soya protein concentration production, In a special extraction process part of the carbohydrates present in soya beans are extracted and concentrated to viscous liquid product. The dry matter content of soya molasses is around 60 %. Most of the carbohydrates in soya molasses are oligosaccharides (abt. 32%). Around 6 to 6,5 % of the product consist of sucrose. The stachyose level in the product is 3 %.

Soya molasses can be used in all types of feed for ruminants and any level of incorporation in the feed. Stachyose and raffinose cannot be digested by single stomach animals, for instance pigs and poultry. Soya molasses should therefore not be used in feeds for poultry (broilers and laying hens) and in feeds for piglets younger than 12 weeks.

In feeds for grower and finisher pigs soya molasses may be used at levels of up to 2 % of the total feed. For pelleted feed for cattle and horses normally 5 % of soya molasses is used whereas in liquid feed for such animals up to 50% of soya molasses can be added.

### Liability

The information contained herein is based on technical data that Liquid Energy Trading Company Swiss SA (later on called LetCo) believes to be reliable and accurate, but disclaims any warrant or guarantee, express or implied. LetCo reserves the right to change information contained herein without prior notice. Any information included herein taken solely or as a whole, does not suggest in any way the product is of satisfactory quality or is adequate for a particular purpose. The purchasing party is not released from the burden of carrying out his own tests and experiments. Because conditions of use of this product are outside our control, LetCo excludes any legal or other liability for the use of this information or any part of it, whether or not based on LetCos negligence, and therefore LetCo shall not be liable for any damages (special, direct, indirect, consequential damages) relating to the use of this information or the use of the product referred herein, solely or in combination with other products. Consequently all liability on the part of LetCo is excluded. Furthermore, our sales and delivery conditions will apply accordingly.